



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

RULE 5. The outside case may be omitted in all instances when the coffin or casket is transported in hearse or undertaker's wagon.

RULE 6. Every outside case shall bear at least four handles, and when over 5 feet 6 inches in length shall bear six handles.

RULE 7. An approved disinfectant fluid shall contain not less than 5 per cent of formaldehyde gas. The term "embalming" as employed in these rules shall require the injection by licensed embalmers of not less than 10 per cent of the body weight, injected arterially in addition to cavity injection, and 12 hours shall elapse between the time of embalming and the shipment of the body.

SOUTH CAROLINA.

Communicable Diseases—List of Diseases Made Notifiable. (Reg. Bd. of H., July 18, 1913.)

The following diseases were made reportable in South Carolina:

Actinomycosis.	Ophthalmia neonatorum (conjunctivitis of newborn infants).
Anthrax.	Paragonimiasis (endemic hemoptysis).
Cancer.	Paratyphoid fever.
Chicken-pox.	Pellagra.
Cholera, Asiatic (also cholera nostras when Asiatic cholera is present or its importation threatened).	Plague.
Continued fever lasting seven days.	Pneumonia (acute).
Dengue.	Poliomyelitis (acute infectious).
Diphtheria.	Rabies.
Dysentery:	Rocky Mountain spotted, or tick fever.
(a) Amebic.	Scarlet fever.
(b) Bacillary.	Septic sore throat.
Favus.	Smallpox.
German measles.	Tetanus.
Glanders.	Trachoma.
Hookworm disease.	Trichinosis.
Leprosy.	Tuberculosis (all forms, the organ or part affected in each case to be specified).
Malaria.	Typhoid fever.
Measles.	Typhus fever.
Meningitis:	Whooping cough.
(a) Epidemic cerebrospinal.	Yellow fever.
(b) Tuberculous.	
Mumps.	

Milk and Milk Products—Production, Care, and Sale.—(Reg. Bd. of H., Dec. 16, 1913.)

Any person owning or managing a dairy or dairy farm, the products of which are sold, shall conform to each and every rule herein set forth for maintaining and handling all such products in a cleanly and sanitary condition.

Cows.—Cows shall be in a healthy condition, and should be tested for tuberculosis once a year, and those reacting or showing evidence of tuberculosis shall be removed.

Cows shall be kept in a cleanly condition, curried and brushed daily. Bedding shall be fresh, and the temperature of the stable kept comfortable.

Location of stable.—Stable shall be located on ground which is well drained and free from any contaminating surroundings.